

B29

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
15 January 2004 (15.01.2004)

PCT

(10) International Publication Number
WO 2004/005252 A1

(51) International Patent Classification⁷: **C07D 209/52**,
403/12, 417/12, 413/12, 401/12, 407/12, A61K 31/403

(21) International Application Number:
PCT/IB2003/001367

(22) International Filing Date: 11 April 2003 (11.04.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PCT/IB02/02663 8 July 2002 (08.07.2002) IB

(63) Related by continuation (CON) or continuation-in-part
(CIP) to earlier application:
US PCT/IB02/02663 (CIP)
Filed on 8 July 2002 (08.07.2002)

(71) Applicant (for all designated States except US): **RAN-
BAXY LABORATORIES LIMITED** [IN/IN]; 19, Nehru
Place, 110019 New Delhi, Delhi (IN).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SALMAN, Mo-
hammad** [IN/IN]; P - 2/19, DLF City, Phase - II, 122001
Gurgaon, Haryana (IN). **MEHTA, Anita** [IN/US]; 756 Old
Checker Road, Buffalo Grove, IL 60089 (US). **SARMA,
Pakala, Kumara, Savithru** [IN/IN]; 1091, 17B, IF-
FCO Colony, 12201 Gurgaon, Haryana (IN). **SHETTY,**

Shankar, Jayram [IN/IN]; U21/14, DLF Colony, Phase -
III, 122001 Gurgaon, Haryana (IN). **DHARMARAJAN,**
Sankaranarayanan [IN/IN]; Gurera House, 811, Sector
17A, 122001 Gurgaon, Haryana (IN). **KUMAR, Naresh**
[IN/IN]; C-1796, Palam Vihar, 122017 Gurgaon, Haryana
(IN). **SILAMKOTI, Arundutt, Viswanatham** [IN/IN];
97, Doveton Road, 500 010 Bolarium, Secunderabad (IN).
CHUGH, Anita [IN/IN]; RA-36, Inderpuri, 110012 New
Delhi, Delhi (IN).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD,
SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US,
UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: **AZABICYCLO DERIVATIVES AS MUSCARINIC RECEPTOR ANTAGONISTS**

(57) Abstract: This invention generally relates to muscarinic receptor antagonists which are useful, among other uses, for the treat-
ment of various diseases of the respiratory, urinary and gastrointestinal systems mediated through muscarinic receptors. Specifically,
the invention relates to derivatives of azabicyclo compounds, including, for example, 6-substituted azabicyclo[3.1.0] hexanes, and
2,4,6-trisubstituted derivatives. The invention also relates to pharmaceutical compositions containing the compounds and the meth-
ods of treating the diseases mediated through muscarinic receptors.

WO 2004/005252 A1

AZABICYCLO DERIVATIVES AS MUSCARINIC RECEPTOR ANTAGONISTS**FIELD OF THE INVENTION**

This invention generally relates to muscarinic receptor antagonists which are
5 useful, among other uses, for the treatment of various diseases of the respiratory, urinary
and gastrointestinal systems mediated through muscarinic receptors. Specifically, the
invention relates to derivatives of azabicyclo compounds, including, for example, 6-
substituted azabicyclo[3.1.0] hexanes, 2,6- and 4,6-disubstituted derivatives, and 2,4,6-
trisubstituted derivatives, as well as pharmaceutical compositions containing such
10 compounds and methods of treating diseases mediated through muscarinic receptors.

BACKGROUND OF THE INVENTION

Muscarinic receptors as members of the G Protein Coupled Receptors (GPCRs)
are composed of a family of 5 receptor sub-types (M_1 , M_2 , M_3 , M_4 and M_5) and are
15 activated by the neurotransmitter acetylcholine. These receptors are widely distributed on
multiple organs and tissues and are critical to the maintenance of central and peripheral
cholinergic neurotransmission. The regional distribution of these receptor sub-types in the
brain and other organs has been documented. For example, the M_1 subtype is located
primarily in neuronal tissues such as cerebral cortex and autonomic ganglia, the M_2
20 subtype is present mainly in the heart where it mediates cholinergically induced
bradycardia, and the M_3 subtype is located predominantly on smooth muscle and salivary
glands (*Nature*, 323, p.411 (1986); *Science*, 237, p.527 (1987)).

A review in *Current Opinions in Chemical Biology*, 3, p. 426 (1999), as well as in
Trends in Pharmacological Sciences, 22, p. 409 (2001) by Eglen et. al., describes the
25 biological potentials of modulating muscarinic receptor subtypes by ligands in different
disease conditions, such as Alzheimer's Disease, pain, urinary disease condition, chronic
obstructive pulmonary disease, and the like.

A review in *J. Med. Chem.*, 43, p. 4333 (2000), by Felder et. al. describes
therapeutic opportunities for muscarinic receptors in the central nervous system and
30 elaborates on muscarinic receptor structure and function, pharmacology and their
therapeutic uses.

The pharmacological and medical aspects of the muscarinic class of acetylcholine
agonists and antagonists are presented in a review in *Molecules*, 6, p. 142 (2001).

Birdsall et. al. in *Trends in Pharmacological Sciences*, 22, p. 215 (2001) have also summarized the recent developments on the role of different muscarinic receptor subtypes using different muscarinic receptor of knock out mice.

Muscarinic agonists such as muscarine and pilocarpine and antagonists such as atropine have been known for over a century, but little progress has been made in the discovery of receptor subtype-selective compounds, making it difficult to assign specific functions to the individual receptors. Although classical muscarinic antagonists such as atropine are potent bronchodilators, their clinical utility is limited due to high incidence of both peripheral and central adverse effects such as tachycardia, blurred vision, dryness of mouth, constipation, dementia, etc. Subsequent development of the quarterly derivatives of atropine such as ipratropium bromide are better tolerated than parenterally administered options, but most of these are not ideal anti-cholinergic bronchodilators, due to lack of selectivity for muscarinic receptor sub-types, resulting in dose-limiting side-effects such as thirst, nausea, mydriasis and those associated with the heart such as tachycardia mediated by the M₂ receptor.

Annual Review of Pharmacological Toxicol., 41, p. 691 (2001), describes the pharmacology of the lower urinary tract infections. Although anti-muscarinic agents such as oxybutynin and tolterodine that act non-selectively on muscarinic receptors have been used for many years to treat bladder hyperactivity, the clinical effectiveness of these agents has been limited due to the side effects such as dry mouth, blurred vision and constipation. Tolterodine is considered to be generally better tolerated than oxybutynin. (Steers et. al., in *Curr. Opin. Invest. Drugs*, 2, 268; Chapple et. al., in *Urology*, 55, 33; Steers et al., *Adult and Pediatric Urology*, ed. Gillenwatter et al., pp 1220-1325, St. Louis, MO; Mosby. 3rd edition (1996)).

There remains a need for development of new highly selective muscarinic antagonists which can interact with distinct subtypes, thus avoiding the occurrence of adverse effects.

Compounds having antagonistic activity against muscarinic receptors have been described in Japanese patent application Laid Open Number 92921/1994 and 135958/1994; WO 93/16048; U.S. Patent No. 3,176,019; GB 940,540; EP 0325 571; WO 98/29402; EP 0801067; EP 0388054; WO 9109013; U.S. Patent No. 5,281,601. Also, U.S. Patent Nos. 6,174,900, 6,130,232 and 5,948,792; WO 97/45414 are related to 1,4-disubstituted piperidine derivatives; WO 98/05641 describes fluorinated,

1,4-disubstitued piperidine derivatives; WO 93/16018 and WO96/33973 are other references of interest.

A report in *J. Med. Chem.*, **44**, p. 984 (2002), describes cyclohexylmethyl piperidiny1 triphenylpropioamide derivatives as selective M₃ antagonist discriminating
5 against the other receptor subtypes.

SUMMARY OF THE INVENTION


In one aspect, azabicyclo derivatives, including, for example, 6-substituted azabicyclo[3.1.0]hexanes, 2,6- and 4,6-disubstituted derivatives, and 2,4,6-trisubstituted
10 derivatives, are provided as muscarinic receptor antagonists which can be useful as safe and effective therapeutic or prophylactic agents for the treatment of various diseases of the respiratory, urinary and gastrointestinal systems. Also provided are processes for synthesizing such compounds.

In another aspect, pharmaceutical compositions containing such compounds are
15 provided together with acceptable carriers, excipients or diluents which can be useful for the treatment of various diseases of the respiratory, urinary and gastrointestinal systems.

The enantiomers, diastereomers, N-oxides, polymorphs, pharmaceutically acceptable salts and pharmaceutically acceptable solvates of these compounds as well as metabolites having the same type of activity are also provided, as well as pharmaceutical
20 compositions comprising the compounds, their metabolites, enantiomers, diastereomers, N-oxides, polymorphs, solvates or pharmaceutically acceptable salts thereof, in combination with a pharmaceutically acceptable carrier and optionally included excipients.

Other aspects will be set forth in the description which follows, and in part will be
25 apparent from the description or may be learnt by the practice of the invention.

In accordance with one aspect, there are provided compounds having the structure of Formula I:



Formula I

and their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, esters, enantiomers, diastereomers, N-oxides, polymorphs, metabolites, wherein

Ar represents an aryl or a heteroaryl ring having 1-2 hetero atoms selected from the group consisting of oxygen, sulphur and nitrogen atoms, the aryl or heteroaryl rings
5 may be unsubstituted or substituted by one to three substituents independently selected from lower alkyl (C₁-C₄), lower perhalo alkyl (C₁-C₄), cyano, hydroxy, nitro, lower alkoxy (C₁-C₄), lower perhalo alkoxy (C₁-C₄), unsubstituted amino, N-lower alkyl (C₁-C₄) or -aryl amino, amino carbonyl, or N-lower alkyl (C₁-C₄) or -aryl amino carbonyl;

R₁ represents a hydrogen, hydroxy, hydroxy methyl, substituted or unsubstituted
10 amino, alkoxy, carbamoyl or halogen (e.g. fluorine, chlorine, bromine and iodine);

R₂ represents alkyl, C₃-C₇ cycloalkyl ring, a C₃-C₇ cyclo alkenyl ring, an aryl, heterocyclic or a heteroaryl ring having 1 to 2 hetero atoms selected from a group consisting of oxygen, sulphur and nitrogen atoms; the aryl, heteroaryl, heterocyclic or a cycloalkyl ring may be unsubstituted or substituted by one to three substituents
15 independently selected from lower alkyl (C₁-C₄), lower perhalo alkyl (C₁-C₄), cyano, hydroxy, nitro, lower alkoxy, lower alkoxy carbonyl, halogen, lower alkoxy (C₁-C₄), lower perhalo alkoxy (C₁-C₄), unsubstituted amino, N-lower alkyl (C₁-C₄) or -aryl amino, amino carbonyl, or N-lower alkyl (C₁-C₄) or -aryl amino carbonyl;

W represents (CH₂)_p, wherein p represents 0 to 1;

20 X represents an oxygen, sulphur, -NR or no atom, wherein R represents hydrogen or (C₁-6) alkyl;

Y represents CHR₅CO or (CH₂)_q wherein R₅ represents hydrogen or methyl and q represents 0 to 4;


Z represents oxygen, sulphur, or NR₁₀, wherein R₁₀ represents hydrogen, or C₁-6
25 alkyl;

Q represents (CH₂)_n (wherein n represents 0 to 4), CHR₈ (wherein R₈ represents H, OH, C₁-6, alkyl, C₁-6 alkenyl, C₁-6 alkoxy) or CH₂CHR₉ (wherein R₉ represents H, OH, lower alkyl (C₁-C₄) or lower alkoxy (C₁-C₄)); and

R₆ and R₇ are independently selected from H, CH₃, COOH, CONH₂, NH₂, and
30 CH₂NH₂.

In accordance with a second aspect, there are provided compounds having the structure of Formula II (Formula I, when R₆ and R₇ = H) and their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, esters, enantiomers, diastereomers,


N-oxides, polymorphs, or metabolites, wherein Ar, R₁, R₂, W, X, Y, Z and Q are as defined for Formula I.




10

Formula II


In accordance with a third aspect, there are provided compounds having the structure of Formula III (Formula I wherein W is (CH₂)_p where p = 0, X is no atom and Y is (CH₂)_q where q = 0, R₆ = H, R₇ = H) and their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, esters, enantiomers, diastereomers, N-oxides, polymorphs, or metabolites, wherein Ar, R₁, R₂, Z and Q are as defined for Formula I.




Formula III

In accordance with a fourth aspect, there are provided compounds having the structure of Formula IV (Formula I wherein W is (CH₂)_p where p = 0, X is no atom and Y is (CH₂)_q where q = 0, R₆ = H, R₇ = H, R₂ = ) and their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, esters, enantiomers, diastereomers, N-oxides, polymorphs, or metabolites, wherein Ar, R₁, Z and Q are as defined for Formula I and r is 1 to 4.


30



Formula IV

In accordance with a fifth aspect, there are provided compounds having the structure of Formula V (Formula I wherein W is $(CH_2)_p$ where $p = 0$, X is no atom and Y is $(CH_2)_q$ where $q=0$, $R_6 = H$, $R_7 = H$, $R_2 =$ , R_1 is hydroxy, Ar is phenyl), and their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, esters, enantiomers, diastereomers, N-oxides, polymorphs, or metabolites, wherein Z and Q are the same as defined for Formula I, s represents 1 or 2.

10



Formula V

15

In accordance with a sixth aspect, there is provided a method for treatment or prophylaxis of an animal or a human suffering from a disease or disorder of the respiratory, urinary and gastrointestinal systems, wherein the disease or disorder is mediated through muscarinic receptors. The method includes administration of at least one compound having the structure of Formula I.

20

In accordance with a seventh aspect, there is provided a method for treatment or prophylaxis of an animal or a human suffering from a disease or disorder associated with muscarinic receptors, comprising administering to a patient in need thereof, an effective amount of a muscarinic receptor antagonist compound as described above.

25

In accordance with an eighth aspect, there is provided a method for treatment or prophylaxis of an animal or a human suffering from a disease or disorder of the respiratory system such as bronchial asthma, chronic obstructive pulmonary disorders (COPD), pulmonary fibrosis, and the like; urinary system which induce such urinary disorders as urinary incontinence, lower urinary tract symptoms (LUTS), etc.; and gastrointestinal system such as irritable bowel syndrome, obesity, diabetes and gastrointestinal hyperkinesis with compounds as described above, wherein the disease or disorder is associated with muscarinic receptors.


30

In accordance with a ninth aspect, there are provided processes for preparing the compounds as described above.

The compounds described herein exhibit significant potency in terms of their activity, as determined by *in vitro* receptor binding and functional assays and *in vivo* experiments using anaesthetized rabbits. The compounds that were found active *in vitro* were tested *in vivo*. Some of the compounds are potent muscarinic receptor antagonists with high affinity towards M₃ receptors. Therefore, pharmaceutical compositions for the possible treatment for the disease or disorders associated with muscarinic receptors are provided. In addition, the compounds can be administered orally or parenterally.

DETAILED DESCRIPTION OF THE INVENTION

The compounds presented herein may be prepared by methods represented by the following reaction sequences:



The compounds of Formula I may be prepared, for example, by the reaction sequence as shown in Scheme I. The preparation comprises reacting a compound of Formula VII with a compound of Formula VI, wherein

Ar represents an aryl or a heteroaryl ring having 1-2 hetero atoms (such as oxygen, sulphur or nitrogen atoms), where the aryl or heteroaryl rings may be unsubstituted or substituted by one to three substituents independently selected from lower alkyl (C₁-C₄), lower perhalo alkyl (C₁-C₄), cyano, hydroxy, nitro, lower alkoxy (C₁-C₄), lower perhalo alkoxy (C₁-C₄), unsubstituted amino, N-lower alkyl (C₁-C₄) or -aryl amino, amino carbonyl, or N-lower alkyl (C₁-C₄) or -aryl amino carbonyl;

R₁ represents a hydrogen, hydroxy, hydroxy methyl, substituted or unsubstituted amino, alkoxy, carbamoyl or halogen (e.g. fluorine, chlorine, bromine and iodine);

R₂ represents alkyl, C₃-C₇ cycloalkyl ring, a C₃-C₇ cyclo alkenyl ring, an aryl, heterocyclic or a heteroaryl ring having 1 to 2 hetero atoms; the aryl, heteroaryl, heterocyclic or a cycloalkyl ring may be unsubstituted or substituted by one to three substituents independently selected from lower alkyl (C₁-C₄), lower perhalo alkyl (C₁-C₄), cyano, hydroxy, nitro, lower alkoxycarbonyl, halogen, lower alkoxy (C₁-C₄), lower perhalo alkoxy (C₁-C₄), unsubstituted amino, N-lower alkyl (C₁-C₄) or -aryl amino, amino carbonyl, or N-lower alkyl (C₁-C₄) or -aryl amino carbonyl;

W represents (CH₂)_p, wherein p represents 0 to 1;

X represents an oxygen, sulphur, -NR or no atom, wherein R represents hydrogen or (C₁-6) alkyl;

Y represents CHR₅CO or (CH₂)_q wherein R₅ represents hydrogen or methyl and q represents 0 to 4;

Z represents oxygen, sulphur, or NR₁₀, wherein R₁₀ represents hydrogen, C₁-6 or alkyl;

Q represents -(CH₂)_n- (wherein n represents 0 to 4), CHR₈ (wherein R₈ represents H, OH, C₁-6, alkyl, C₁-6 alkenyl, C₁-6 alkoxy) or CH₂CHR₉ (wherein R₉ represents H, OH, lower alkyl (C₁-C₄) or lower alkoxy (C₁-C₄));

R₆ and R₇ are independently selected from H, CH₃, COOH, CONH₂, NH₂, and CH₂NH₂; and

P is any protecting group for an amino group, for example, benzyl or t-butyloxy carbonyl groups.

The reaction between a compound of Formula VII and a compound of Formula VI can take place in the presence of a condensing agent (for example, 1-(3-

dimethylaminopropyl)-3-ethyl carbodiimide hydrochloride (EDC) or 1,8-diazabicyclo
[5.4.0]undec-7-ene (DBU)), in a solvent (such as N,N-dimethylformamide,
dimethylsulfoxide, toluene, or xylene, at temperatures ranging from about 0 to about
140°C), to give a protected compound of Formula VIII which on deprotection in the
5 presence of a deprotecting agent (for example, palladium on carbon, trifluoroacetic acid
(TFA) or hydrochloric acid) in an organic solvent (for example, methanol, ethanol,
tetrahydrofuran or acetonitrile, at temperatures ranging from about 10 to about 50°C)
gives an unprotected compound of Formula I.

In the above scheme, where specific bases, condensing agents, protecting groups,
10 deprotecting agents, solvents, catalysts, temperatures, etc. are mentioned, it is to be
understood that other bases, condensing agents, protecting groups, deprotecting agents,
solvents, catalysts, temperatures, etc. known to those skilled in the art may be used.
Similarly, the reaction temperature and duration may be adjusted according to the desired
needs.


15 Suitable salts of the compounds represented by the Formula I were prepared so as
to solubilize the compound in aqueous medium for biological evaluations, as well as to be
compatible with various dosage formulations and also to aid in the bioavailability of the
compounds. Examples of such salts include pharmacologically acceptable salts such as
inorganic acid salts (for example, hydrochloride, hydrobromide, sulphate, nitrate and
20 phosphate), organic acid salts (for example, acetate, tartarate, citrate, fumarate, maleate,
tolounesulphonate and methanesulphonate). When carboxyl groups are included in the
Formula I as substituents, they may be present in the form of an alkaline or alkali metal
salt (for example, sodium, potassium, calcium, magnesium, and the like). These salts
may be prepared by various techniques, such as treating the compound with an equivalent
25 amount of inorganic or organic, acid or base in a suitable solvent.

Particular compounds are shown here:

- (2R,2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-
cyclopentyl-2-phenyl acetamide (Compound 1);
(2R,2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-
30 cyclopentyl-2-phenyl acetamide hydrochloride salt (Compound 2);
(2R)-(1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-
cyclopentyl 2-phenyl acetamide (Compound 3);
(2R)-(1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-
cyclopentyl 2-phenyl acetamide hydrochloride salt (Compound 4);

- (2S)-(1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl 2-phenyl acetamide (Compound 5);
- (2S)-(1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl 2-phenyl acetamide hydrochloride salt (Compound 6);
- 5 (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-methoxy-2-cyclopentyl-2-phenyl acetamide (Compound 7);
- (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cycloheptyl-2-phenyl acetamide (Compound 8);
- (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclobutyl-2-phenyl acetamide (Compound 9);
- 10 (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclobutyl-2-phenyl acetamide tartarate salt (Compound 10);
- (2R) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-(3,3-difluorocyclopentyl)-2-phenyl acetamide (Compound 11);
- 15 (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-(3-fluorocyclopentyl)-2-phenyl acetamide (Compound 12);
- (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-(3,3-difluorocyclopentyl)-2-phenyl acetamide (Compound 13);
- (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-(3,3-difluorocyclopentyl)-2-phenyl acetamide tartarate salt (Compound 14);
- 20 (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2,2-diphenyl acetate (Compound 15);
- (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2,2-diphenyl acetamide (Compound 16);
- 25 (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclohexyl-2-phenyl acetamide (Compound 17) and
- (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hex-6-yl methyl]-2-cyclopentyl-2-hydroxy-N-methyl-2-phenyl acetamide (Compound 18).

Table I



5

Formula III

(Formula I, wherein $W=(CH_2)_p$ where $p=0$, X is no atom and $Y=(CH_2)_q$, where $q=0$, $R_6=R_7=H$)

Compound No.	Ar	R ₁	R ₂	Z	Q
1. (2R,2S)	-C ₆ H ₅	-OH	cyclopentyl	NH-	-CH ₂ -
2. (HCl salt) (2R,2S)	-C ₆ H ₅	-OH	cyclopentyl	NH-	-CH ₂ -
3. (2R)	-C ₆ H ₅	-OH	cyclopentyl	NH-	-CH ₂ -
4. (HCl salt) (2R)	-C ₆ H ₅	-OH	cyclopentyl	NH-	-CH ₂ -
5. (2S)	-C ₆ H ₅	-OH	cyclopentyl	NH-	-CH ₂ -
6. (HCl salt) (2S)	-C ₆ H ₅	-OH	cyclopentyl	NH-	-CH ₂ -
7. (2R, 2S)	-C ₆ H ₅	-OCH ₃	cyclopentyl	NH-	-CH ₂ -
8. (2R, 2S)	-C ₆ H ₅	-OH	cycloheptyl	NH-	-CH ₂ -
9. (2R, 2S)	-C ₆ H ₅	-OH	cyclobutyl	NH-	-CH ₂ -
10.	-C ₆ H ₅	-OH	cyclobutyl	NH-	-CH ₂ -

Compound No.	Ar	R ₁	R ₂	Z	Q
(2R, 2S) Tartarate salt					
11. (2R)	-C ₆ H ₅	-OH	3,3-difluoro cyclopentyl	NH-	-CH ₂ -
12. (2R, 2S)	-C ₆ H ₅	-OH	3-fluoro cyclopentyl	NH-	-CH ₂ -
13. (2R, 2S)	-C ₆ H ₅	-OH	3,3-difluoro cyclopentyl	NH-	-CH ₂ -
14. (2R, 2S) Tartarate salt	-C ₆ H ₅	-OH	3,3-difluoro cyclopentyl	NH-	-CH ₂ -
15. (2R, 2S)	-C ₆ H ₅	-OH	phenyl	O	-CH ₂ -
16. (2R, 2S)	-C ₆ H ₅	-OH	phenyl	NH-	-CH ₂ -
17. (2R, 2S)	-C ₆ H ₅	-OH	cyclohexyl	NH-	-CH ₂ -
18. (2R, 2S)	-C ₆ H ₅	-OH	cyclopentyl	-N-CH ₃	-CH ₂ -
19.	<i>p</i> -F-C ₆ H ₄	-F	-C ₆ H ₅	NH-	-CH(OH)-
20.	<i>o</i> -Cl- <i>p</i> -F- C ₆ H ₃	-NH ₂	1-cyclohexenyl	-O-	-CH(OH)-
21.	3-pyrrolyl	-Cl	<i>p</i> -CH ₃ -C ₆ H ₄	-O-	- CH(CH ₃)-
22.	3-pyridyl	CH ₂ (OH)	Cyclobutyl	-O-	-CH(OH)-
23.	3-furyl	-F	-C ₆ H ₅	N(CH ₃)	-CH ₂ CH ₂ -
24.	1-indolyl	-Cl	Cyclohexyl	-S-	-CH ₂ -
25.	3-isothiazolyl	-F	-C ₆ H ₅	-NH-	-CH(OH)-
26.	3-isoxazolyl	N(CH ₃) ₂	<i>o</i> -Cl- <i>p</i> -F-C ₆ H ₃	-O-	-CH ₂ CH ₂ -

Because of their valuable pharmacological properties, the compounds described herein may be administered to an animal for treatment orally, or by a parenteral route. The pharmaceutical compositions described herein can be produced and administered in dosage units, each unit containing a certain amount of at least one compound described
5 herein and/or at least one physiologically acceptable addition salt thereof. The dosage may be varied over extremely wide limits as the compounds are effective at low dosage levels and relatively free of toxicity. The compounds may be administered in the low micromolar concentration, which is therapeutically effective, and the dosage may be increased as desired up to the maximum dosage tolerated by the patient.

10 The compounds described herein can be produced and formulated as their enantiomers, diastereomers, N-Oxides, polymorphs, solvates and pharmaceutically acceptable salts, as well as metabolites having the same type of activity. Pharmaceutical compositions comprising the molecules of Formulae I, II, III, IV and V or metabolites, enantiomers, diastereomers, N-oxides, polymorphs, solvates or pharmaceutically
15 acceptable salts thereof, in combination with pharmaceutically acceptable carrier and optionally included excipient can also be produced.

Compounds such as those described in Table I, for example, can be prepared from the appropriate acetic acid, in analogy to specific examples provided below. The preparation of such acetic acids is known to those of ordinary skill in the art, and will be
20 recognized by reference to the cited specific examples given below. Other compounds within the scope of the invention, such as those having R_6 and/or R_7 as methyl, carboxylic acid, amide, amino or methylamino will be readily prepared in analogy to the specific procedures given in the specific examples below, using appropriate azabicyclo[3.1.0]hexanes which are synthesized using known methods. Other compounds
25 within the scope of the invention, such as those having X as oxygen, sulfur or secondary or tertiary amine can be prepared in analogy to the specific procedures given in the specific examples below, using appropriate esters, thiocompounds, or amides which can be made by procedures known to those of ordinary skill in the art. Similarly, compounds within the scope of the invention, such as those having Y as CHR_5CO where R_5 is
30 hydrogen or methyl can be made in analogy to the specific procedures given in the specific examples below, using appropriate anhydrides, imides or thioanhydrides which are made by procedures known to those of skill in the art. Other compounds within the scope of the invention, such as those having Z as oxygen, sulfur or secondary or tertiary amino can be made in analogy to the specific procedures given in the specific examples

below, using appropriate starting materials which are made by procedures known to those of ordinary skill in the art.

The examples mentioned below demonstrate general synthetic procedures, as well as specific preparations of particular compounds. The examples are provided to illustrate the details of the invention and should not be constrained to limit the scope of the present invention.

EXAMPLES

Various solvents, such as acetone, methanol, pyridine, ether, tetrahydrofuran, hexanes, and dichloromethane, were dried using various drying reagents according to procedures described in the literature. IR spectra were recorded as nujol mulls or a thin neat film on a Perkin Elmer Paragon instrument, Nuclear Magnetic Resonance (NMR) were recorded on a Varian XL-300 MHz instrument using tetramethylsilane as an internal standard.

Example 1: Preparation of (2R,2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide (Compound No.1)

Step a: Synthesis of (2R,2S) (1 α ,5 α ,6 α)-N-[3-benzyl 3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide

To a solution of (1 α ,5 α ,6 α)-6-aminomethyl-3-benzyl-3-azabicyclo[3.1.0]hexane (prepared as described in EP 0413455 A2) (29.9 mmol, 6.05 gm) in dimethyl formamide (100 ml) was added 2-(R,S)-hydroxy-2-cyclopentyl-2-phenyl acetic acid (prepared following *J. Amer Chem. Soc.*, 1953; 75:2654) (27.2 mmol, 6.0 gm) and cooled to 0°C. The reaction mixture was treated with hydroxy benzotriazole (29.9 mmol, 4.04 gm) followed by N-methyl morpholine (54.4 mmol, 5.2 gm) was stirred at 0°C for 0.5 hours. EDC (1-[3-(dimethylamino)propyl]-3-ethyl carbodiimide hydrochloride (29.9mmol, 5.7 gms) was added and the reaction mixture was stirred at 0° C for 1 hour and further at room temperature (RT) overnight. The reaction mixture was poured into saturated sodium bicarbonate and extracted with ethyl acetate. The organic layers were washed with water and dried over sodium sulphate and concentrated under reduced pressure. The residue was purified by column chromatography (silica gel 100-200 mesh) eluting compound in 93-95% purity. To obtain higher purity (about 99%) of the compound, it was triturated with toluene and filtered.

Step b: Preparation of (2R,2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide

To a solution of (2R,2S) (1 α ,5 α ,6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-aminomethyl-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide from Example 1, step a
5 (1.0 g, 2.48 mmol) in methanol (25.0 ml), 5% Pd-C (0.2 g), (50% wet) was added under N₂. Then, anhydrous ammonium formate (0.8 g, 12.38 mmol) was added under stirring and the reaction mixture was refluxed for half an hour under the N₂ atmosphere. The mixture was cooled to room temperature and the reaction mixture was filtered through a bed of hyflo. The hyflo bed was washed with methanol (75.0 ml), ethyl acetate (25.0 ml)
10 and water (25.0 ml). The filtrate was concentrated under vacuum. The residue was diluted with water and pH of the resulting solution was adjusted to pH~14 with 1N NaOH. The solution was extracted with ethyl acetate (2x50 ml) and the ethyl acetate layer was washed with water and brine solution. The layer was dried over anhydrous Na₂SO₄ and concentrated to give the title compound as solid in 96.2% (0.75g, 2.39 mmol)
15 yield with >98 % purity by HPLC.

The compound exhibited a melting point of 149-151°C, and had infrared absorbance (KBr) at 3410, 2951.5, 2868.3, and 1652.5 cm⁻¹. ¹HNMR (CDCl₃) spectral data were as follows: δ 7.59-7.62 (m, 2H), 7.23-7.36 (m, 3H), 6.56 (brs, 1H), 3.03-3.15 (m, 3H), 2.78-2.90 (m, 4H, including OH), 1.51-1.71 (m, 8H), 1.19-1.27 (m, 4H), 0.70-
20 0.72 (m, 1H). The mass spectrum showed peaks at m/e of 315 (MH⁺), 297 (M-OH).

Example 2: Preparation of (2R,2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide. Hydrochloride salt (Compound No.2)

To a solution of (2R,2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide (prepared in Example 1)
25 (0.2g, 0.637 mmole) in dichloromethane (4.0 ml), ethanolic HCL (1.45 N, 0.5 ml, 0.725 mmol) was added at room temperature and stirred for 10 minutes. At the same temperature diethyl ether (100 ml) was added to the reaction mixture, stirred for 5
30 minutes and concentrated under vacuum without heating. The residue was triturated with ether to get solid material. The ether layer was decanted and the solid was dried under vacuum to get the title compound as a hygroscopic solid in 94% (0.21g, 0.6 mmole) yield with >98% purity by HPLC.

Example 3: Preparation of (2R)-(1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide (Compound No.3)

5 **Step a: Synthesis of (2R)-(1 α ,5 α ,6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide**

This compound was synthesised following the procedure of Example 1, Step a, using (2R)-2-hydroxy-2-cyclopentyl-2-phenyl acetic acid (synthesised as in Grover *et. al.*, *J. Org. Chem.*, 2000; 65:6283-6287), instead of 2-hydroxy-2-cyclopentyl-2-phenyl acetic acid.

10 **Step b: Synthesis of (2R)-(1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide**

This compound was synthesised following the procedure of Example 1, Step b, using (2R)- (1 α ,5 α ,6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide instead of (2R,2S) (1 α ,5 α ,6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl
15 acetamide. The enantiomeric excess (ee) was determined by HPLC (Chinacel OD, mobile phase 90% hexane/10% EtOH/0.1% TFA) by observing the (S) and (R) isomers. The (S) isomer eluted at approximately 11.11 min. The (R) isomer eluted at approximately 11.81 min. The optical purity was > 99%.

20 The compound exhibited a melting point of 150.2⁰C. Infrared spectral data showed (DCM): 1653.8 cm⁻¹. ¹HNMR spectral data showed (CDCl₃): δ 7.61 (d, J=9Hz, 2H), 7.30-7.38 (m, 2H), 6.70 (s, 1H), 3.61-3.68 (m, 2H), 3.08-3.28 (m, 5H), 1.49-1.68 (m, 10H), 1.11-1.26 (m, 2H), 0.75-0.85 (m, 1H).

25 Example 4: Preparation of (2R)-(1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide. hydrochloride salt (Compound No.4)

The hydrochloride salt was synthesised following the same procedure as in Example 2, using (2R)-(1 α , 5 α , 6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-
30 hydroxy-2-cyclopentyl-2-phenyl acetamide instead of (2R,2S) (1 α , 5 α , 6 α)-N-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide.

Example 5: Preparation of (2S)-(1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide (Compound No.5)

Step a: Synthesis of (2S)-(1 α ,5 α ,6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide

5 This compound was synthesized following the procedure of Example 1, Step a, using (2S)-2-hydroxy-2-cyclopentyl-2-phenyl acetic acid (synthesised as in Grover *et. al.*, *J. Org. Chem.*, 2000; 65:6283-6287), instead of 2-hydroxy-2-cyclopentyl-2-phenyl acetic acid.

Step b: Synthesis of (2S)-(1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide

10 This compound was synthesized following the procedure of Example 1, Step b, using (2S)-(1 α ,5 α ,6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide instead of (2R,2S) (1 α ,5 α ,6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl
15 acetamide. The ee was determined by HPLC (Chinacel OD, mobile phase 90% hexane/10% EtOH/0.1% TFA) by observing the (S) and (R) isomers. The (S) isomer eluted at approximately 11.11 min. The (R) isomer eluted at approximately 11.81 min. The optical purity was > 99%.

The compound exhibited a melting point of 62.6-63.3 °C. Infrared spectral data
20 showed (KBr): 1653.7 cm⁻¹. ¹HNMR spectral data showed (CDCl₃): δ 7.59-7.62 (m, 2H), 7.29-7.37 (m, 3H), 3.58-3.65 (m, 2H), 3.02-3.24 (m, 4H), 1.11-1.34 (m, 11H), 0.75-0.95 (m, 1H).

Example 6: Preparation of (2S)-(1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide. hydrochloride salt (Compound No.6)

25 The hydrochloride salt was synthesized in 90 % yield following the procedure of Example 2, using (2S)-(1 α , 5 α , 6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide instead of (2R,2S) (1 α , 5 α , 6 α)-N-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl
30 acetamide.

Example 7: Preparation of (2R, 2S) (1 α , 5 α , 6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-methoxy-2-cyclopentyl-2-phenyl acetamide (Compound 7)

5 Step a: Preparation of (2R, 2S) 2-methoxy-2-cyclopentyl-2-phenyl acetic acid ethyl ester

To a cold solution of (2R, 2S) 2-hydroxy-2-cyclopentyl-2-phenyl acetic acid ethyl ester (synthesized as per *J. Am. Chem. Soc.*, 1953; 75:2654) (4.5 mmol) in dimethyl formamide was added sodium hydride (9.08 mmol) in portions at 0°C and stirred at room
10 temperature for 1 hr. Reaction mixture was cooled to 0°C and iodomethane (18.0 mmol) was added. Reaction mixture was then stirred at room temperature for 2 hrs. TLC showed absence of starting material. Water was added to reaction mixture extracted with ethylacetate. The organic layer dried over anhydrous sodium sulfate and concentrated. Crude compound was purified by column chromatography & desired product eluted with
15 2% EtOAc/Hexane.

¹H NMR (CDCl₃) spectral data were as follows: δ 7.47-7.36 (5H, m), 4.31 (2H, q), 3.26 (3H, s), 2.43 (1H, m), 1.66-1.46 (11H, m)

Step b: Synthesis of (2R, 2S) 2-methoxy-2-cyclopentyl-2-phenyl acetic acid

To a solution of (2R, 2S) 2-methoxy-2-cyclopentyl-2-phenyl acetic acid ethyl
20 ester (1.8 mmol) in methanol, potassium hydroxide (KOH) (2.2 mmol) was added and the reaction mixture refluxed for 7 hrs. TLC showed presence of starting material then 3 mole equivalent KOH was added and reaction mixture refluxed for 3 hrs. TLC showed absence of starting materials. Reaction mixture was concentrated, residue was taken in water and neutralized with concentrated hydrochloric acid & extracted with ethylacetate.
25 Organic layer was washed with water, brine, dried over anhydrous sodium sulphate and concentrated under reduced pressure to obtain the desired compound.

¹H NMR (CDCl₃) spectral data were as follows: δ 7.48-7.35 (5H, m), 3.20 (3H, s), 2.94-2.86 (1H, m), 1.86-1.50 (8H, m)

30 Step c: Preparation of (2R, 2S) (1 α , 5 α , 6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-methoxy-2-cyclopentyl-2-phenyl acetamide

This was prepared following the procedure of Example 1, step 'a' by using 2-methoxy-2-cyclopentyl-2-phenyl acetic acid instead of 2-hydroxy-2-cyclopentyl-2-phenyl acetic acid.

Step d: Preparation of (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-methoxy-2-cyclopentyl-2-phenyl acetamide

This was prepared following the procedure of Example 1, step b by using (2R, 2S) (1 α , 5 α , 6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-methoxy-2-cyclopentyl-2-phenyl acetamide instead of (2R, 2S) (1 α , 5 α , 6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide

¹H NMR (CDCl₃) spectral data showed: δ 7.45-7.30 (5H, m), 7.03 (1H, m), 3.25-3.02 (9H, m), 2.00-0.86 (12H, m).

10

Example 8: Preparation of (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cycloheptyl-2-phenyl acetamide (Compound 8)

Step a: Synthesis of (2R, 2S)-(1 α ,5 α ,6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cycloheptyl-2-phenyl acetamide

This compound was synthesized following the procedure of Example 1, step a, using (2R, 2S)-2-hydroxy-2-cycloheptyl-2-phenyl acetic acid (synthesized as in Grover *et. al.*, *J. Org. Chem.*, 2000; 65:6283-6287), instead of 2-hydroxy-2-cyclopentyl-2-phenyl acetic acid.

Step b: Synthesis of (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cycloheptyl-2-phenyl acetamide

This compound was synthesized following the procedure of Example 1, step b, using (2R, 2S)-(1 α -5 α , 6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cycloheptyl-2-phenyl acetamide instead of (2R, 2S) (1 α , 5 α , 6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide in 90% yield.

The compound had

¹H NMR spectral data showed (CDCl₃): δ 7.59-7.61 (m, 2H), 7.13-7.36 (m, 3H), 6.76 (brs, 1H), 3.00-3.20 (m, 2H), 2.80-2.92 (m, 2H), 2.50-2.80 (m, 1H), 2.40 (brs, 2H), 1.28-1.73 (m, 12H), 1.00-1.20 (m, 2H), 0.80-0.90 (m, 1H).

Infrared absorbance (DCM) at 1655.7 cm⁻¹

The mass spectrum showed peaks at m/e of : 343 (MH⁺)

Example 9: Preparation of (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclobutyl-2-phenyl acetamide (Compound 9)

Step a: synthesis of (2R, 2S) (1 α ,5 α ,6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclobutyl-2-phenyl acetamide

This compound was synthesized following the procedure of Example 1, step a, using (2R, 2S) 2-hydroxy-2-cyclobutyl-2-phenyl acetic acid (synthesized as per reported
5 procedure of Saul B. Kadin and Joseph G.Cannon., *J. Org. Chem.*, 1962; 27:240-245), instead of 2-hydroxy-2-cyclopentyl-2-phenyl acetic acid.

Step b: Synthesis of (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclobutyl-2-phenyl acetamide

This compound was synthesized following the procedure of Example 1, step b,
10 using (2R, 2S) (1 α , 5 α , 6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclobutyl-2-phenyl acetamide instead of (2R, 2S) (1 α , 5 α , 6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide to give the title compound with 90.6% purity by HPLC.

¹H NMR (CDCl₃) spectral data were as follows: δ 7.50 (2H, m), 7.30 (3H, m), 6.60 (1H,
15 m), 3.60-3.00 (9H, m), 2.04 (1H, m), 1.96-1.74 (6H, m), 1.45 (1H, m), 1.00 (1H, m).

Example 10: Preparation of (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclobutyl-2-phenyl acetamide tartarate salt (Compound 10)

20 To a solution of compound No. 9 in ethanol, solid tartaric acid was added and the solution was stirred for 1 hour at room temperature and the solvent was evaporated. Ether was added to it to precipitate the salt. It was washed with ether (4 times) by decanting supernatant liquid to give the salt as powder with 95.66% purity by HPLC.

¹H NMR (CDCl₃) spectral data were as follows: δ 7.45 (2H, m), 7.21 (3H, m), 4.36 (2H,
25 s), 3.42 (2H, m), 3.20 (3H, m), 3.05 (2H, m), 1.97 (1H, m), 1.90-1.60 (7H, m), 1.10 (1H, m).

Example 11: Preparation of (2R) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-(3,3-difluoro cyclopentyl)-2-phenyl acetamide (Compound 11)

30 **Step a: Preparation of (2R,5R)-2-tert-butyl-5-phenyl-1,3-dioxalan-4-one**

The compound was synthesised following the procedure described in *J. Org. Chem.*, 2000; 65:6283-6287.

Step b: Preparation of (2R,5R)-2-tert-butyl-5-[(1Ror1S)-3-oxocyclopentyl]-5-phenyl-1,3-dioxalan-4-one

To a suspension of compound of step a (1.36mmol) in tetrahydrofuran (THF) (12ml) was added lithium diisopropyl amide (LDA) in THF (1.5 mmol) drop wise at –
5 78°C under nitrogen atmosphere. The reaction mixture was stirred at the same temperature for 2 hours. A solution of 2-cyclopenten-1-one (1.52mmol) in THF (2ml) was added to the reaction mixture dropwise and stirred for additional 3hours. The reaction mixture was quenched with saturated aq.ammonium chloride solution and extracted with ethyl acetate. The organic layer was dried and the residue obtained after removing the
10 solvents in vacuo was purified by column chromatography (100-200mesh silica gel). The product was eluted with 10% EtOAc–hexane mixture.

¹HNMR(CDCl₃):δ-values: 7.70-7.26 (m,5Ar-H), 5.43-5.37 (d,1H), 2.91-2.88 (m,1H), 2.37-1.77 (m,6H), 0.92 (s, 9H)

IR(DCM): 1791 and 1746 cm⁻¹

Step c: Preparation of (2R, 5R)-2-tert-butyl-5-[(1Ror1S)-3,3-difluorocyclopentyl]-5-phenyl-1,3-dioxalan-4-one

To a solution of compound of step-b (1mmol) in chloroform (15ml) was added diethylamino sulphurtrifluoride (DAST), (3.3 mmol) at 0°C under nitrogen atmosphere. The reaction mixture was stirred at the same temperature for 30 minutes and then at room
20 temperature for 3 days. After being cooled to 0°C, the reaction mixture (RM) was quenched carefully by adding water. The organic layer was separated and the aqueous layer extracted with chloroform. The combined organic layers were dried and the residue obtained after removing the solvent was purified by column chromatography (100-200 mesh size silica gel) eluting the compound with 5% EtOAc-hexane mixture.

25 ¹HNMR(CDCl₃):δ-values : 7.73-7.35 (m,5Ar-H), 5.49 (s,1H), 2.86-2.82 (m,1H), 2.27-1.80 (m,6H), 0.98 (s,H)

IR(DCM): 1793 cm⁻¹

Step d: Preparation of (2R)- [(1Sor1R)-3,3-difluorocyclopentyl]-2- hydroxy-2-phenylaceticacid

30 The solution of compound of step-c (1mmol) in MeOH (10ml) was stirred with 3N aqueous sodium hydroxide solution for overnight at room temperature. The reaction mixture was concentrated under reduced pressure. The residue was diluted with water and extracted with dichloromethane. The aqueous layer was acidified with conc.hydrochloric

acid and extracted with EtOAc. The organic layer was dried and concentrated under reduced pressure to give the product.

m.pt. :123°C

¹HNMR(CDCl₃):δ-values : 7.69-7.37(m,5Ar-H) , 3.29-3.20(m,1H) , 2.39-1.68 (m,6H)

5 **Step e: Preparation of (1α,5α,6α)-6-aminomethyl-3-benzyl-3-azabicyclo[3.1.0]hexane.**

The compound was synthesised as per the procedure of EP 0413455A2.

Step f: Preparation of (2R)- (1α,5α,6α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-[(1R or 1S)-3,3-difluorocyclopentyl]-2- hydroxy-2-phenylacetamide.

It was prepared following the procedure of Example 1, step a, using the acid synthesized in step d instead of 2-hydroxy-2-cyclopentyl-2-phenyl acetic acid.

Step g: Synthesis of (2R) (1α,5α,6α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-(3,3-difluorocyclopentyl)-2-phenyl acetamide

15 It was prepared by following the procedure of Example 1, step b by using (2R) (1α, 5α, 6α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-(3,3-difluorocyclopentyl)-2-phenyl acetamide instead of (2R, 2S) (1α, 5α, 6α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide. The optical purity is 87.27% (HPLC).

20 ¹H NMR (CDCl₃) spectral data showed: δ 7.59-7.55 (2H, m), 7.35-7.31 (3H, m), 7.03 (1H, m), 3.18-3.11 (7H, m), 1.87-1.62 (9H, m).

Example 12: Preparation of (2R, 2S) (1α,5α,6α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-(3-fluorocyclopentyl)-2-phenyl acetamide (Compound 12)

25 **Step-a: Preparation of (2R,2S)-2-tert-butyl-5-[(1R or 1S, 3R or 3S)-3-hydroxy cyclopentyl]-5-phenyl-1,3-dioxalan-4-one.**

 To a solution of (2R, 2S, 5R)-2-tert-butyl-5-[(1R or 1S)-3-oxocyclopentyl]-5-phenyl-1,3-dioxalan-4-one (1 mmol) in methanol (10 ml) cooled to 0°C, sodium borohydride (2 mmol) was added in small lots with stirring. The RM was stirred at 0°C
30 for 1 hr. It was concentrated under reduced pressure and the residue diluted with water and extracted with EtOAc. The organic layer was dried and the residue obtained after the

removal of solvents was purified by column chromatography (100-200 mesh silica gel) eluting the compound with 20% EtOAc-hexane mixture.

$^1\text{H NMR}(\text{CDCl}_3)$: δ -values : 7.68-7.29 (m, 5H, ArH) , 5.45 (d, 1H) , 4.30 (m, 1H), 3.25 (m, 1H), 2.65-2.63 (m, 1H), 1.80-1.63 (m, 6H), 0.92 (s, 9H)

5 IR(DCM): 1789 cm^{-1} , 3386 cm^{-1}

Step-b: Preparation of (2R, 2S)-2-tert-butyl-5-[1R or 1S, 3R or 3S]-3-fluorocyclopentyl]-5-phenyl-1,3-dioxolan-4-one.

The solution of compound of step-a (1 mmol) in chloroform (10 ml) was cooled to 0°C and diethylamino sulphur trifluoride (DAST) (1.5 mmol) was added dropwise under nitrogen atmosphere. The RM was stirred at 0°C for 30 minutes and then at room temperature for 3 days. The RM was cooled and carefully quenched with aqueous ammonium chloride solution. The organic layer was separated and aqueous layer extracted with EtOAc. The combined organic layer was dried and residue obtained after removing the solvents was purified by column chromatography (100-200 mesh, silica gel) eluting the compound with 5% EtOAc-hexane mixture.

$^1\text{H NMR}(\text{CDCl}_3)$: δ -values : 7.68-7.28 (m, 5H, Ar-H) , 5.46 (d, 1H), 5.39 (m, 1H), 2.90 (m, 1H), 1.98-1.25 (m, 6H), 0.93 (s, 9H)

Step-c: Preparation of (2R, 2S)-[(1R or 1S, 3R or 3S)-3-fluorocyclopentyl]-2-hydroxy-2-phenylacetic acid.

20 The compound was synthesised following the procedure of Example 11, step-d using (2R, 2S, 5R)-2-tert-butyl-5-[(1R or 1S, 3R or 3S)-3-fluorocyclopentyl]-5-phenyl-1,3-dioxolan-4-one instead of (2R, 5R)-2-tert-butyl-5-[(1R or 1S)-3,3-difluorocyclopentyl]-5-phenyl-1,3-dioxolan-4-one.

$^1\text{H NMR}(\text{CDCl}_3)$: δ 7.66-7.27 (m, 5Ar-H), 5.30-5.00 (m, H), 3.32-3.16 (m, 1H), 2.05-1.26 (m, 6H).

25 IR(DCM): 1710 cm^{-1}

Step-d: Preparation of (2R, 2S)-(1 α ,5 α ,6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-[1R or 1S, 3R or 3S]-3-fluorocyclopentyl]-2-hydroxy-2-phenylacetamide.

30 This compound was synthesized following the procedure of Example 1, step a, using the acid synthesized in the above step-c, instead of 2-hydroxy-2-cyclopentyl-2-phenyl acetic acid.

Step-e: Preparation of (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-(3-fluorocyclopentyl)-2-phenyl acetamide

This compound was synthesized following the procedure of Example 1, step b, using (2R, 2S)-(1 α ,5 α ,6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-[1R or 1S, 3R or 3S]-3-fluorocyclopentyl]-2-hydroxy-2-phenylacetamide instead of (2R, 2S) (1 α , 5 α , 6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide. The optical purity was 87.27% (HPLC).

¹H NMR spectral data showed (CDCl₃): δ 7.56 (2H, m), 7.35 (3H, m), 6.08 (1H, m), 5.30-5.03 (1H, m), 3.27 (1H, m), 3.11 (2H, m), 2.91 (4H, m), 2.04-1.48 (9H, m), 0.71 (1H, m).

10

Example 13: Preparation of (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-(3,3-difluorocyclopentyl)-2-phenyl acetamide (Compound 13)

This compound was prepared following the procedure of Example 11, by using (2R, 2S) [(1 α , 5 α , 6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-(3,3-difluorocyclopentyl)-2-phenyl acetamide] instead of (2R) [(1 α , 5 α , 6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-(3,3-difluorocyclopentyl)-2-phenyl acetamide] in step g, Example 11.

The optical purity is 83.778% (HPLC).

¹H NMR spectral data showed (CDCl₃) δ : 7.57-7.30 (5H, m), 6.49-6.44 (1H, m), 3.33 (1H, m), 3.10 (2H, m), 6.49-6.44 (1H, m), 3.33 (1H, m), 3.10 (2H, m), 2.87 (3H, m), 2.23-1.80 (8H, m), 1.79-1.20 (2H, m)

IR (KBr) : 3410, 1654 cm⁻¹

Example 14: Preparation of (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-(3,3-difluorocyclopentyl)-2-phenyl acetamide tartarate salt (Compound 14)

To a solution of compound No. 13 in ethanol was added tartaric acid & heated the reaction mixture at 60°C for 1 hour. The reaction mixture was then concentrated under reduced pressure, added diethyl ether and the organic layer was removed to get light brown solid as the desired compound. The optical purity was found to be 98.14% (HPLC)

¹H NMR spectral data showed (CD₃OD): δ 7.50 (2H, m), 7.20 (3, m), 4.40 (2H, s), 3.60-3.00 (6H, m), 2.10-1.60 (8H, m), 1.19 (1H, m), 0.90 (1H, m).

IR (KBr) : 3407, 1653 cm^{-1}

Example 15: Preparation of (2R, 2S) (1 α , 5 α , 6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(methyl)-yl]-2-hydroxy-2,2-diphenyl acetate (Compound 15)

5 **Step a: Synthesis of (2R, 2S) (1 α , 5 α , 6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(methyl)-yl]-2-hydroxy-2,2-diphenyl acetate.**

Step (i): Preparation of (1 α , 5 α , 6 α)-3-benzyl-6-hydroxymethyl-3-azabicyclo[3.1.0]hexane.

 Synthesized as per reported procedure of EP 0 413 455 A2

10 **Step (ii): Preparation of (1 α , 5 α , 6 α)-3-benzyl-6-(methanesulfonyloxy)methyl-3-azabicyclo[3.1.0]hexane:**

 A solution of the title compound of preparation of step (i) (0.203 g; 1 mmol) and triethyl amine (0.21 gms, 2 mmol) in ethyl acetate (25 ml) was cooled to -10°C and treated with methanesulfonyl chloride (0.17 gms, 1.5 mmol). After stirring for one hour
15 at -10°C , the reaction was poured into a saturated aqueous sodium bicarbonate solution. The organic layer was dried over sodium sulphate. Filtration and removal of solvent *in vacuo* provided the title compound as a yellow oil, which was used as such in the following step without further purification.

^1H NMR (CDCl_3): δ -values: 7.45 (m, 5H, arom.), 4.29 (s, 2H), 3.81 (m, 2H), 3.13 (m,
20 4H), 2.84 (s, 3H), 1.38 (m, 3H)

Step (iii): Preparation of (1 α , 5 α , 6 α)-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(methyl)-yl]-2-hydroxy-2,2-diphenyl acetate

 To a solution of 2-hydroxy-2,2-diphenyl acetic acid (synthesized as per reported procedures in Vogel's textbook of "Practical Organic Chemistry," page 1046 (5th Ed); *J.*
25 *Am. Chem. Soc.*, 1953; 75:2654 and EP 613232) (1 mmol, 0.228 gms) in xylene was added, (1 α , 5 α , 6 α)-3-benzyl-6-(methanesulfonyloxy) methyl-3-azabicyclo[3.1.0]hexane (0.28 gms, 1 mmol) followed by DBU (1,8-diazabicyclo[5,4,0] undec-7-ene, (2 mmol, 0.305 gms) and the reaction mixture refluxed for 6 hrs. The RM was then washed with water, brine and dried over sodium sulphate. The solvents were evaporated and the crude
30 compound thus obtained was purified by column chromatography (silicagel, 100-200 mesh) eluting the compound with 20-80, ethylacetate hexane.

^1H NMR (CDCl_3): δ - values: 7.46-7.22 (m, 15H, arom), 4.24 (s, 1H), 4.11-4.09 (d, 2H), 3.56 (s, 2H), 2.91-2.89 (d, 2H), 2.31-2.29 (d, 2H), 1.67-1.62 (m, 1H), 1.3 (s, 2H)

IR (DCM): 1724 cm^{-1}

Step b: Synthesis of (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(methyl)-yl]-2-hydroxy-2,2-diphenyl acetate

It was debenzylated by following the procedure of Example 1, step b to give the
5 title compound in 60% yield.

IR (KBr): 1731.6 cm^{-1}

^1H NMR (CDCl_3) spectral data showed: δ 7.33-7.47 (m, 10H), 4.17 (d, 2H, $J=6\text{Hz}$), 2.72-2.92 (m, 4H), 0.94-0.99 (m, 2H), 0.88 (t, 1H),

The mass spectrum showed peaks at m/e of 324 ($M+1$)

10

Example 16: Preparation of (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2,2-diphenyl acetamide (Compound 16)

Step a: Preparation of (2R, 2S) (1 α ,5 α ,6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2,2-diphenyl acetamide.

15 It was prepared following the procedure of Example 1, step a by using 2-hydroxy-2,2-diphenyl acetic acid (synthesized as per reported procedures in Vogel's textbook of "Practical Organic Chemistry," page 1046 (5th Ed), *J. Am. Chem. Soc.*, 1953; 75:2654 and EP 613232) instead of 2-hydroxy-2-cyclopentyl-2-phenyl acetic acid.

Step b: Preparation of (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-(3,3-difluorocyclopentyl)-2-phenyl acetamide

20 This compound was prepared following the procedure of Example 1, step b, using (1 α ,5 α ,6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2,2-diphenylacetamide instead of (2R, 2S) (1 α ,5 α ,6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide to give the title
25 compound in 70% yield.

IR (KBr): 1658.0 cm^{-1}

^1H NMR (CDCl_3) spectral data showed: δ 7.34-7.44 (m, 10H), 6.53 (s, 1H), 3.17-3.26 (m, 2H), 2.87-3.01 (m, 4H), 1.38 (s, 2H), 0.88 (t, 1H).

The mass spectra showed peaks at m/e 323 ($M+1$), 305 ($M-\text{OH}$).

30

Example 17: Preparation of (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclohexyl-2-phenyl acetamide (Compound 17)

Step a: Synthesis of (1 α ,5 α ,6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclohexyl-2-phenyl acetamide

It was prepared following the procedure of Example 1, step a by using 2-hydroxy-2-cyclohexyl-2-phenyl acetic acid (synthesized as per the procedure described in *J. Amer. Chem. Soc.*, 1953; 75:2654) instead of 2-hydroxy-2-cyclopentyl-2-phenyl acetic acid.

Step b: Preparation of (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-cyclohexyl-2-phenyl acetamide

This compound was prepared following the procedure of Example 1, step b using (2R, 2S) (1 α ,5 α ,6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclohexyl-2-phenyl acetamide instead of (2R, 2S) (1 α ,5 α ,6 α)-N-[3-benzyl-3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide to give the title compound in 80% yield.

IR (KBr): 1654.7 cm⁻¹

¹H NMR (CDCl₃) spectral data showed: δ 7.59-7.62 (m, 2H), 7.29-7.37 (m, 3H), 6.71 (s, 1H), 3.03-3.14 (m, 2H), 2.80-2.92 (m, 4H), 2.42 (m, 1H), 1.13-1.35 (m, 12H), 0.88 (m, 1H)

The mass spectra recorded peaks of 329 (M+1), 311 (M-OH).

Example 18: Preparation of (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hex-6-yl methyl]-2-cyclopentyl-2-hydroxy-N-methyl-2-phenyl acetamide (Compound 18)

Step a: Preparation of (2R, 2S) (1 α ,5 α ,6 α)-N-[3-tert-butyloxycarbonyl-3-azabicyclo[3.1.0]-hexyl-6-yl methyl]-2-hydroxy-2-cyclopentyl-2-phenyl acetamide

To a solution of (1 α ,5 α ,6 α)-3N-benzyl-6-amino-3-azabicyclo[3.1.0] hexane (synthesized as per reported procedure of Braish T.F. et. al., *Synlett*, 1996; 1100) (2.5 g, 7.96 mole) in (50.0 mmol) at 0°C, triethylamine (3.9 ml, 28 mmol) and Boc-anhydride (5.2 g, 23.9 mmol) were added. The reaction mixture was stirred at 0°C for 30 minute and at room temperature for 12 hrs. Diluted with dichloromethane (50 ml) and washed with water and brine solution. Dried over anhydrous sodium sulfate and concentrated. The residue was purified by column chromatography using 25% ethyl acetate in hexane to give the title compound as solid in 86% (2.85 g, 6.9 mmol) yield. The compound exhibited a melting point of 179.5-182.9°C.

Step b: Preparation of (2R, 2S) (1 α ,5 α ,6 α)-N-[3-tert-butyloxycarbonyl-3-azabicyclo[3.1.0]hexyl-6-yl methyl]-2-(3-silyltrimethyloxy)-2-cyclopentyl-2-phenyl acetamide

To a solution of Boc-derivative (2.0 g, 4.8 mmol) in dimethylformamide (10.0 ml), imidazole (1.2 g, 16.9 mmol) and trimethylsilyl chloride (1.54 ml, 12.0 mmol) were added and the reaction mixture was stirred at RT for 12 hrs. Diluted with water (50.0 ml) and extracted with ethyl acetate. The ethyl acetate layer was washed with water and brine solution. Dried and concentrated. The residue was purified by column chromatography using 15% ethyl acetate in hexane to give the title compound in 85% (2.0 g, 4.1 mmol) yield.

Step c: Preparation of (2R, 2S) (1 α ,5 α ,6 α)-N-[3-tert-butyloxycarbonyl-3-azabicyclo[3.1.0]hex-6-yl methyl)-2-cyclopentyl-2-hydroxy-N-methyl-2-phenyl acetamide

To a solution of trimethylsilyl derivative (2.0 g, 4.1 mmol) and n-tetrabutyl ammonium iodide (0.11 gm, 0.3 mmol) in dry tetrahydrofuran (THF) (20.9 ml) at 0°C, sodium hydride (0.6 g, 12.3 mmol) was added in portion and the resulting solution was stirred at 0°C for 15 minutes and allowed to cool to room temperature and stirred for 1 hours at RT. Again cooled to 0°C and methyl iodide (2.3 ml, 36.8 mmol) in dry THF (2.0 ml) was added dropwise. Stirred for 12 hrs at RT. Saturated aqueous NaHCO₃ solution (10.0 ml) was added organic layer was separated and dried over anhydrous sodium sulfate. Concentrated and the residue was purified by column chromatograph using 15% ethyl acetate in hexane to give the title compound as semisolid in 61% (1.25 g, 2.49 mmol) yield.

Step d: Preparation of (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hex-6-ylmethyl)-2-cyclopentyl-2-hydroxy-N-methyl-2-phenyl acetamide

To a solution of the compound (0.2 g, 0.4 mmol) from the above step in ethanol (5.0 ml), conc.HCl was added dropwise till the pH of the reaction mixture was 2. The reaction mixture was stirred at RT for 24 hrs. Neutralized with saturated aqueous sodium bicarbonate solution. Concentrated under vacuum and the residue was taken in dichloromethane (10.0 ml) and washed with water and brine solution. The organic layer was dried over anhydrous sodium sulfate and concentrated to give the title compound as solid in 54% (0.07 g, 0.21 mmol) yield.

m.p.: 91.5°C

¹H NMR (CDCl₃) spectral data showed: δ 7.29-7.42 m, (5H), 5.39 (m, 1H), 2.81-3.52 (m, 10H), 1.11-1.82 (m, 12H)

IR (DCM): 1621.9 cm^{-1}

The mass spectrum showed peak of 329 (MH^+).

Biological Activity

5 **Radioligand Binding Assays:** The affinity of test compounds for M_2 and M_3 muscarinic receptor subtypes was determined by [^3H]-N-methylscopolamine binding studies, using rat heart and submandibular gland, respectively, as described by Moriya et al., (*Life Sci.*, 1999; 64(25):2351-2358) with minor modifications as follows. The membrane preparation was done with the following modifications: a low spin step of
10 500g for 10 minutes at 4°C was used; the buffer was 20 mM HEPES, 10 mM EDTA, at pH 7.4; the high speed spin was done at 40,000g and the homogenate was passed through a filter gauge before any spinning. The assay conditions were modified as follows: the assay volume was 250 μL ; the incubation time was 3 hours; the PE concentration was 0.1%; the filtermat used was GF/B from Wallac; the scintillant used was Supermix from
15 Wallac; the amount of scintillant was 500 μL /well; and the counter used was a 1450 microbeta PLUS, from Wallac.

Membrane preparation: Submandibular glands and heart were isolated and placed in ice cold homogenising buffer (HEPES 20mM, 10mM EDTA, pH 7.4) immediately after sacrifice. The tissues were homogenised in 10 volumes of
20 homogenising buffer and the homogenate was filtered through two layers of wet gauze and filtrate was centrifuged at 500g for 10min. The supernatant was subsequently centrifuged at 40,000g for 20 min. The pellet thus obtained was resuspended in same volume of assay buffer (HEPES 20 mM, EDTA 5mM, pH 7.4) and were stored at -70°C until the time of assay.

25 **Ligand binding assay:** The compounds were dissolved and diluted in DMSO. The membrane homogenates (150-250 μg protein) were incubated in 250 μl of assay buffer (HEPES 20 mM, pH 7.4) at 24-25°C for 3h. Non-specific binding was determined in the presence of 1 μM atropine. The incubation was terminated by vacuum filtration over GF/B fiber filters (Wallac). The filters were then washed with ice cold 50mM Tris
30 HCl buffer (pH 7.4). The filter mats were dried and bound radioactivity retained on filters was counted. The IC_{50} and K_d were estimated by using the non-linear curve fitting program using G Pad Prism software. The value of inhibition constant K_i was calculated from competitive binding studies by using Cheng & Prusoff equation (*Biochem*

Pharmacol, 1973; 22:3099-3108), $K_i = IC_{50} / (1 + L/K_d)$, where L is the concentration of [3H]NMS used in the particular experiment. $pK_i = -[\log K_i]$

Functional Experiments using isolated rat bladder:

5 **Methodology:** Animals were euthanized by overdose of urethane and whole bladder was isolated and removed rapidly and placed in ice cold Tyrode buffer with the following composition (mMol/L) NaCl 137; KCl 2.7; CaCl₂ 1.8; MgCl₂ 0.1; NaHCO₃ 11.9; NaH₂PO₄ 0.4; glucose 5.55 and continuously gassed with 95% O₂ and 5 % CO₂.

10 The bladder was cut into longitudinal strips (3mm wide and 5-6 mm long) and mounted in 10 ml organ baths at 30° C, with one end connected to the base of the tissue holder and the other end connected to a polygraph through a force displacement transducer. Each tissue was maintained at a constant basal tension of 2 g and allowed to equilibrate for 1 hour during which the PSS was changed every 15 min. At the end of equilibration period the stabilization of the tissue contractile response was assessed with
15 1 µmol/L of Carbachol consecutively for 2-3 times. Subsequently a cumulative concentration response curve to carbachol (10^{-9} mol/L to 3×10^{-5} mol/L) was obtained. After several washes, once the baseline was achieved, cumulative concentration response curve was obtained in presence of NCE (NCE added 20 min. prior to the second CRC).

20 The contractile results were expressed as % of control E max. ED₅₀ values were calculated by fitting a non-linear regression curve (Graph Pad Prism). pKB values were calculated by the formula $pKB = -\log [(\text{molar concentration of antagonist} / (\text{dose ratio} - 1))]$ where, dose ratio = ED₅₀ in the presence of antagonist/ED₅₀ in the absence of antagonist.

25 ***In vivo* experiments using anesthetized rabbit:** The effect of test substances was studied on carbachol evoked changes on bladder pressure, heart rate and salivation. Male rabbits weighing 1.2-3 kg were anaesthetized with urethane (1.5g/kg), and administered as a slow intravenous infusion through the marginal ear vein. The tracheae were cannulated to maintain airway patency. Blood pressure was recorded from the femoral artery by means of a Statham P10 EZ pressure transducer connected to a Grass
30 model 7D polygraph. The heart rate was monitored by a tachograph triggered by the pulse wave of blood pressure. The other femoral artery was cannulated for the administration of carbachol. Test compound and saline were infused intravenously via the femoral vein.

The bladder was exposed through a midline laparotomy and both the ureters were identified, carefully separated and ligated. The ureters were incised proximally to allow free flow of urine from the kidney to the exterior. Bladder neck was gently held and the urethra was traced and separated from the adjoining tissues. PE canula was introduced
5 into the bladder and ligated. The bladder was drained and subsequently filled with 15ml of warm saline (37°C). The other end of the intravesical catheter was connected to the Grass model 7D polygraph through a Statham P10 EZ pressure transducer to monitor the bladder pressure. Care was taken to keep the exposed area moist and warm. A period of 30-60 min was allowed for stabilization of parameters subsequent to surgery. Salivation
10 response was assessed by placing preweighed absorbent cotton gauze in the buccal cavity for 2 minutes after carbachol administration.

The effect of the compound on carbachol (1.5 µg/kg, intrarterial) induced changes on blood pressure, heart rate and bladder pressure were observed. At least two stable responses were obtained. These responses were considered as 100%. Subsequently,
15 effect of increasing dose of test compound or vehicle (i.v, 12 to 15 min before carbachol challenge) was studied.

The change in bladder pressure, salivation and agonist induced bradycardia were expressed as % change from pretreatment control. ID₅₀ values (dose required to inhibit 50% of response) were calculated from non-linear curve fitting for sigmoidal dose
20 response curve using Graph Pad Prism software and values were expressed as µg/kg . The results of the *in-vitro* and *in-vivo* tests are listed in Table II and Table III.

Table II

Compd.No.	Receptor Binding Assay		Functional Assay pK _B
	M ₃ Pki	M ₂ pKi	
1	9.28	7.92	9.36± 0.17
2	9.5± 0.03	8.36± 0.03	9.16± 0.18
4	9.65± 0.03	8.43± 0.07	9.28± 0.2
5	7.5± 0.04	6.8± 0.04	-
6	7.88± 0.03	7.11± 0.07	7.8± 0.09
7	8.61	7.6	8.56± 0.11
8	8.83	7.35	-
9	8.9	8.02	-
10	9.5	8.2	8.73± 0.17
11	8.9	7.4	-
12	9.04	7.6	-
13	8.7	7.3	-
14	8.87	7.14	8.73± 0.18
15	10.18	9.66	-
16	9.41	7.95	-
17	8.87	7.3	-
18	9.42	8.56	-
Tolterodine	8.47	8.68	8.86± 0.12


5

Table III

Compound No.	ID ₅₀ (µg/kg,i.v)			Fold Selectivity	
	Bladder Pressure	Salivary Response	Heart Rate Response	Bladder <i>versus</i> Salivary	Bladder <i>versus</i> Heart Rate
Tolterodine	25.84 ± 4.24	32.5 ± 10.02	32.54 ± 5.77	1.24± 0.21	1.26± 0.19
2	3.93 ± 1.1	7.67 ± 1.19	21.13 ± 4.47	3.08 ± 0.94	12.6 ± 5.7
4	2.14 ± 0.42	4.66 ± 0.56	5.65 ± 0.86	2.37 ± 0.22	3.43 ± 1.02

WE CLAIM:

1. Compounds having the structure of Formula I:

**Formula - I**

and their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, esters, enantiomers, diastereomers, N-oxides, polymorphs, or metabolites, wherein

Ar represents an aryl or a heteroaryl ring having 1-2 hetero atoms, the aryl or heteroaryl rings may be unsubstituted or substituted by one to three substituents independently selected from lower alkyl (C₁-C₄), lower perhalo alkyl (C₁-C₄), cyano, hydroxy, nitro, lower alkoxy (C₁-C₄), lower perhalo alkoxy (C₁-C₄), unsubstituted amino, N-lower alkyl (C₁-C₄) or -aryl amino, amino carbonyl, or N-lower alkyl (C₁-C₄) or -aryl amino carbonyl;

R₁ represents a hydrogen, hydroxy, hydroxy methyl, substituted or unsubstituted amino, alkoxy, carbamoyl or halogen;

R₂ represents alkyl, C₃-C₇ cycloalkyl ring, a C₃-C₇ cyclo alkenyl ring, an aryl, heterocyclic or a heteroaryl ring having 1 to 2 hetero atoms; the aryl, heteroaryl, heterocyclic or a cycloalkyl ring may be unsubstituted or substituted by one to three substituents independently selected from lower alkyl (C₁-C₄), lower perhalo alkyl (C₁-C₄), cyano, hydroxy, nitro, lower alkoxy, carbonyl, halogen, lower alkoxy (C₁-C₄), lower perhalo alkoxy (C₁-C₄), unsubstituted amino, N-lower alkyl (C₁-C₄) or -aryl amino, amino carbonyl, or N-lower alkyl (C₁-C₄) or -aryl amino carbonyl;

W represents (CH₂)_p, wherein p represents 0 to 1;

X represents an oxygen, sulphur, -NR or no atom, wherein R represents hydrogen or (C₁-6) alkyl;


Y represents CHR₅CO or (CH₂)_q wherein R₅ represents hydrogen or methyl and q represents 0 to 4;

Z represents oxygen, sulphur, or NR₁₀, wherein R₁₀ represents hydrogen or C₁-6 alkyl;

32 Q represents $-(CH_2)_n-$, wherein n represents 0 to 4, CHR_8 , wherein R_8 represents
 33 H, OH, C_{1-6} , alkyl, C_{1-6} alkenyl, or C_{1-6} alkoxy, or Q represents CH_2CHR_9 , wherein
 34 R_9 represents H, OH, lower alkyl (C_1-C_4) or lower alkoxy (C_1-C_4); and


35 R_6 and R_7 are independently selected from H, CH_3 , $COOH$, $CONH_2$, NH_2 , and
 36 CH_2NH_2 .

- 1 2. The compounds according to claim 1 having the structure of Formula II and their
 2 pharmaceutically acceptable salts, pharmaceutically acceptable solvates, esters,
 3 enantiomers, diastereomers, N-oxides, polymorphs, or metabolites.




9 Formula II

- 1 3. The compounds according to claim 1 having the structure of Formula III and their
 2 pharmaceutically acceptable salts, pharmaceutically acceptable solvates, esters,
 3 enantiomers, diastereomers, N-oxides, polymorphs, or metabolites.




9 Formula III

- 1 4. The compounds according to claim 1 having the structure of Formula IV and their
 2 pharmaceutically acceptable salts, pharmaceutically acceptable solvates, esters,
 3 enantiomers, diastereomers, N-oxides, polymorphs, or metabolites, wherein r is 1 to 4.



9 Formula IV


- 1 5. The compounds according to claim 1 having the structure of Formula V, and their
 2 pharmaceutically acceptable salts, esters, enantiomers, N-oxides, or metabolites;
 3 wherein s represents 1 to 2.



10 **Formula V**

- 1 6. A compound selected from the group consisting of
- 2 (2R,2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-
 3 cyclopentyl-2-phenyl acetamide (Compound 1);
- 4 (2R,2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-
 5 cyclopentyl-2-phenyl acetamide hydrochloride salt (Compound 2);
- 6 (2R)-(1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-
 7 cyclopentyl 2-phenyl acetamide (Compound 3);
- 8 (2R)-(1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-
 9 cyclopentyl 2-phenyl acetamide hydrochloride salt (Compound 4);
- 10 (2S)-(1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-
 11 cyclopentyl 2-phenyl acetamide (Compound 5);
- 12 (2S)-(1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-
 13 cyclopentyl 2-phenyl acetamide hydrochloride salt (Compound 6);
- 14 (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-methoxy-2-
 15 cyclopentyl-2-phenyl acetamide (Compound 7);
- 16 (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-
 17 cycloheptyl-2-phenyl acetamide (Compound 8);
- 18 (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-
 19 cyclobutyl-2-phenyl acetamide (Compound 9);
- 20 (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-
 21 cyclobutyl-2-phenyl acetamide tartarate salt (Compound 10);
- 22 (2R) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-(3,3-
 23 difluorocyclopentyl)-2-phenyl acetamide (Compound 11);
- 24 (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-
 25 (3-fluorocyclopentyl)-2-phenyl acetamide (Compound 12);
- 26 (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-
 27 (3,3-difluorocyclopentyl)-2-phenyl acetamide (Compound 13);

- 28 (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-
 29 (3,3-difluorocyclopentyl)-2-phenyl acetamide tartarate salt (Compound 14);
 30 (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-
 31 2,2-diphenyl acetate (Compound 15);
 32 (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-
 33 2,2-diphenyl acetamide (Compound 16);
 34 (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hexyl-6-(aminomethyl)-yl]-2-hydroxy-2-
 35 cyclohexyl-2-phenyl acetamide (Compound 17) and
 36 (2R, 2S) (1 α ,5 α ,6 α)-N-[3-azabicyclo[3.1.0]hex-6-ylmethyl)-2-cyclopentyl-2-
 37 hydroxy-N-methyl-2-phenyl acetamide (Compound 18).
 38 7. A pharmaceutical composition comprising a therapeutically effective amount of a
 39 compound as defined in claim 1, 2, 3, 4, 5 or 6 together with pharmaceutically
 40 acceptable carriers, excipients or diluents.
 1 8. A method for treatment or prophylaxis of an animal or a human suffering from a
 2 disease or disorder of the respiratory, urinary and gastrointestinal systems, wherein
 3 the disease or disorder is mediated through muscarinic receptors, comprising
 4 administering to said animal or human, a therapeutically effective amount of a
 5 compound having the structure of Formula I,



Formula I

12 its pharmaceutically acceptable salts, pharmaceutically acceptable solvates, esters,
 13 enantiomers, diastereomers, N-oxides, polymorphs or metabolites, wherein

14 Ar represents an aryl or a heteroaryl ring having 1-2 hetero atoms, wherein the
 15 aryl or heteroaryl rings may be unsubstituted or substituted by one to three
 16 substituents independently selected from lower alkyl (C₁-C₄), lower perhalo alkyl (C₁-
 17 C₄), cyano, hydroxy, nitro, lower alkoxy (C₁-C₄), lower perhalo alkoxy (C₁-C₄),
 18 unsubstituted amino, N-lower alkyl (C₁-C₄) or -aryl amino, amino carbonyl, or N-
 19 lower alkyl (C₁-C₄) or -aryl amino carbonyl;

20 R₁ represents a hydrogen, hydroxy, hydroxy methyl, amino, alkoxy, carbamoyl or
 21 halogen;

R₂ represents alkyl, C₃-C₇ cycloalkyl ring, a C₃-C₇ cyclo alkenyl ring, an aryl, heterocyclic or a heteroaryl ring having 1 to 2 hetero atoms; the aryl, heteroaryl, heterocyclic or a cycloalkyl ring may be unsubstituted or substituted by one to three substituents independently selected from lower alkyl (C₁-C₄), lower perhalo alkyl (C₁-C₄), cyano, hydroxy, nitro, lower alkoxy, carbonyl, halogen, lower alkoxy (C₁-C₄), lower perhalo alkoxy (C₁-C₄), unsubstituted amino, N-lower alkyl (C₁-C₄) or -aryl amino, amino carbonyl, or N-lower alkyl (C₁-C₄) or -aryl amino carbonyl;

W represents (CH₂)_p, wherein p represents 0 to 1;

X represents an oxygen, sulphur, -NR or no atom, wherein R represents hydrogen or (C₁-6) alkyl;


Y represents CHR₅CO or (CH₂)_q wherein R₅ represents hydrogen or methyl and q represents 0 to 4;

Z represents oxygen, sulphur, or NR₁₀, wherein R₁₀ represents hydrogen or C₁-6 alkyl;

Q represents -(CH₂)_n-, wherein n represents 0 to 4, CHR₈, wherein R₈ represents H, OH, C₁-6, alkyl, C₁-6 alkenyl, C₁-6 alkoxy, or Q represents CH₂CHR₉, wherein R₉ represents H, OH, lower alkyl (C₁-C₄) or lower alkoxy (C₁-C₄); and


R₆ and R₇ are independently selected from H, CH₃, COOH, CONH₂, NH₂, CH₂NH₂.

9. The method according to claim 8 for treatment or prophylaxis of an animal or a human suffering from a disease or disorder of the respiratory, urinary and gastrointestinal systems, wherein the disease or disorder is mediated through muscarinic receptors, comprising administering to said animal or human, a therapeutically effective amount of a compound having the structure of Formula II, its pharmaceutically acceptable salts, pharmaceutically acceptable solvates, esters, enantiomers, diastereomers, N-oxides, polymorphs or metabolites.




Formula II

- 1 10. The method according to claim 8 for treatment or prophylaxis of an animal or a
 2 human suffering from a disease or disorder of the respiratory, urinary and
 3 gastrointestinal systems, wherein the disease or disorder is mediated through
 4 muscarinic receptors, comprising administering to said animal or human, a
 5 therapeutically effective amount of a compound having the structure of Formula III,
 6 its pharmaceutically acceptable salts, pharmaceutically acceptable solvates, esters,
 7 enantiomers, diastereomers, N-oxides, polymorphs, prodrugs or metabolites.



8
 9
 10
 11
 12 **Formula III**


- 1 11. The method according to claim 8 for treatment or prophylaxis of an animal or a
 2 human suffering from a disease or disorder of the respiratory, urinary and
 3 gastrointestinal systems, wherein the disease or disorder is mediated through
 4 muscarinic receptors, comprising administering to the said animal or human, a
 5 therapeutically effective amount of a compound having the structure of Formula IV,
 6 its pharmaceutically acceptable salts, pharmaceutically acceptable solvates, esters,
 7 enantiomers, diastereomers, N-oxides, polymorphs, prodrugs or metabolites, wherein
 8 r is 1 to 4.



9
 10
 11
 12
 13
 14
 15 **Formula IV**

- 1 12. The method according to claim 8 for treatment or prophylaxis of an animal or a
 2 human suffering from a disease or disorder of the respiratory, urinary and
 3 gastrointestinal systems, wherein the disease or disorder is mediated through
 4 muscarinic receptors, comprising administering to said animal or human, at least one
 5 therapeutically effective amount of a compound having the structure of Formula V, its
 6 pharmaceutically acceptable salts, pharmaceutically acceptable solvates, esters,

enantiomers, diastereomers, N-oxides, polymorphs or metabolites, wherein s represents 1 to 2.




Formula V

13. The method according to claim 8 wherein the disease or disorder is urinary incontinence, lower urinary tract symptoms (LUTS), bronchial asthma, chronic obstructive pulmonary disorders (COPD), pulmonary fibrosis, irritable bowel syndrome, obesity, diabetes or gastrointestinal hyperkinesis.
14. The method according to claim 9 wherein the disease or disorder is urinary incontinence, lower urinary tract symptoms (LUTS), bronchial asthma, chronic obstructive pulmonary disorders (COPD), pulmonary fibrosis, irritable bowel syndrome, obesity, diabetes or gastrointestinal hyperkinesis.
15. The method according to claim 10 wherein the disease or disorder is urinary incontinence, lower urinary tract symptoms (LUTS), bronchial asthma, chronic obstructive pulmonary disorders (COPD), pulmonary fibrosis, irritable bowel syndrome, obesity, diabetes or gastrointestinal hyperkinesis.
16. The method according to claim 11 wherein the disease or disorder is urinary incontinence, lower urinary tract symptoms (LUTS), bronchial asthma, chronic obstructive pulmonary disorders (COPD), pulmonary fibrosis, irritable bowel syndrome, obesity, diabetes or gastrointestinal hyperkinesis.
17. The method according to claim 12 wherein the disease or disorder is urinary incontinence, lower urinary tract symptoms (LUTS), bronchial asthma, chronic obstructive pulmonary disorders (COPD), pulmonary fibrosis, irritable bowel syndrome, obesity, diabetes or gastrointestinal.
18. The method for treatment or prophylaxis of an animal or a human suffering from a disease or disorder of the respiratory, urinary and gastrointestinal systems, wherein the disease or disorder is mediated through muscarinic receptors, comprising

administering to said animal or human, a therapeutically effective amount of the pharmaceutical composition according to claim 7.

19. The method according to claim 18 wherein the disease or disorder urinary incontinence, lower urinary tract symptoms (LUTS), bronchial asthma, chronic obstructive pulmonary disorders (COPD), pulmonary fibrosis, irritable bowel syndrome, obesity, diabetes or gastrointestinal hyperkinesis.

20. A method of preparing a compound of Formula I,



Formula I

and its pharmaceutically acceptable salts, pharmaceutically acceptable solvates, esters, enantiomers, diastereomers, N-oxides, polymorphs or metabolites, wherein

Ar represents an aryl or a heteroaryl ring having 1-2 hetero atoms, wherein the aryl or heteroaryl rings may be unsubstituted or substituted by one to three substituents independently selected from lower alkyl (C₁-C₄), lower perhalo alkyl (C₁-C₄), cyano, hydroxy, nitro, lower alkoxy (C₁-C₄), lower perhalo alkoxy (C₁-C₄), unsubstituted amino, N-lower alkyl (C₁-C₄) or -aryl amino, amino carbonyl, or N-lower alkyl (C₁-C₄) or -aryl amino carbonyl;

R₁ represents a hydrogen, hydroxy, hydroxy methyl, amino, alkoxy, carbamoyl or halogen;

R₂ represents alkyl, C₃-C₇ cycloalkyl ring, a C₃-C₇ cyclo alkenyl ring, an aryl, heterocyclic or a heteroaryl ring having 1 to 2 hetero atoms selected from a group consisting of oxygen, sulphur and nitrogen atoms; the aryl, heteroaryl, heterocyclic or a cycloalkyl ring may be unsubstituted or substituted by one to three substituents independently selected from lower alkyl (C₁-C₄), lower perhalo alkyl (C₁-C₄), cyano, hydroxy, nitro, lower alkoxy carbonyl, halogen, lower alkoxy (C₁-C₄), lower perhalo alkoxy (C₁-C₄), unsubstituted amino, N-lower alkyl (C₁-C₄) or -aryl amino, amino carbonyl, or N-lower alkyl (C₁-C₄) or -aryl amino carbonyl;

W represents (CH₂)_p, wherein p represents 0 to 1;

X represents an oxygen, sulphur, -NR or no atom, wherein R represents hydrogen or (C₁-6) alkyl;

Y represents CHR_5CO or $(\text{CH}_2)_q$ wherein R_5 represents hydrogen or methyl and q represents 0 to 4;


Z represents oxygen, sulphur, NR_{10} , wherein R_{10} represents hydrogen, C_{1-6} alkyl;

Q represents $(\text{CH}_2)_n$ (wherein n represents 0 to 4), CHR_8 (wherein R_8 represents H, OH, C_{1-6} alkyl, C_{1-6} alkenyl, C_{1-6} alkoxy) or CH_2CHR_9 (wherein R_9 represents H, OH, lower alkyl ($\text{C}_1\text{-C}_4$) or lower alkoxy ($\text{C}_1\text{-C}_4$)); and


R_6 and R_7 are independently selected from H, CH_3 , COOH , CONH_2 , NH_2 , CH_2NH_2 ;

said method comprising:

(a) reacting a compound of Formula VII with a compound of Formula VI




Formula VII




Formula VI

to give a protected compound of Formula VIII wherein Ar, R_1 , R_2 , W, X, Y, Z, and Q are as defined, and P is a protecting group for an amino group



Formula VIII

(b) deprotecting the compound of Formula VIII in the presence of a deprotecting agent to give compound of Formula I wherein Ar, R_1 , R_2 , W, X, Y, Z, and Q are as defined,



Formula I

21. The method of claim 20, wherein P is any protecting group for an amino group and is selected from the group consisting of benzyl and t-butyloxy carbonyl groups.

- 1 22. The method of claim 20, wherein the reaction of a compound of Formula VI with a
2 compound of Formula VII to give a compound of Formula VIII is carried out in the
3 presence of a condensing agent which is selected from the group consisting of 1-(3-
4 dimethyl amino propyl)-3-ethyl carbodiimide hydrochloride (EDC) and 1,8-
5 diazabicyclo [5.4.0]undec-7-ene (DBU).
- 1 23. The method of claim 20, wherein the reaction of a compound of Formula VI with a
2 compound of Formula VII is carried out in a suitable polar aprotic solvent selected
3 from the group consisting of N,N-dimethylformamide, dimethyl sulfoxide, toluene,
4 and xylene.
- 1 24. The method of claim 20, wherein the reaction of compound of Formula VI with a
2 compound of Formula VII is carried out at 0-140°C.
- 1 25. The method of claim 20, wherein the deprotection of a compound of Formula VIII is
2 carried out with a deprotecting agent which is selected from the group consisting of
3 palladium on carbon, trifluoroacetic acid (TFA) and hydrochloric acid.
- 1 26. The method of claim 20, wherein the deprotection of a compound of Formula VIII to
2 give a compound of Formula I is carried out in a suitable organic solvent selected
3 from the group consisting of methanol, ethanol, tetrahydrofuran and acetonitrile.

INTERNATIONAL SEARCH REPORT

Internat Application No
PCT/IB 03/01367

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D209/52 C07D403/12 C07D417/12 C07D413/12 C07D401/12
C07D407/12 A61K31/403

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 863 141 A (BANYU PHARMA CO LTD) 9 September 1998 (1998-09-09) page 3, line 29 -page 3, line 55	1-26
A	WO 02 051841 A (ALMIRALL PRODESFARMA SA ;BUIL ALBERO MARIA ANTONIA (ES); FERNANDEZ) 4 July 2002 (2002-07-04) page 3, line 22 -page 5, line 23	1-26
A	WO 02 04402 A (BANYU PHARMA CO LTD ;MATSUDA KENJI (JP); KURIHARA HIDEKI (JP); OGI) 17 January 2002 (2002-01-17) claim 1	1-26

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *g* document member of the same patent family

Date of the actual completion of the international search

21 July 2003

Date of mailing of the international search report

28/07/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Uselli, A

INTERNATIONAL SEARCH REPORT

Int'l application No.
PCT/IB 03/01367**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 8-19 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compounds.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest☐ The additional search fees were accompanied by the applicant's protest.☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Internat	Application No
PCT/LB 03/01367	

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0863141	A	09-09-1998	AT 205490 T	15-09-2001
			AU 7145996 A	30-04-1997
			DE 69615214 D1	18-10-2001
			DE 69615214 T2	27-06-2002
			EP 0863141 A1	09-09-1998
			US 6130232 A	10-10-2000
			CA 2234619 A1	17-04-1997
			WO 9713766 A1	17-04-1997
<hr/>				
WO 02051841	A	04-07-2002	WO 02051841 A1	04-07-2002
<hr/>				
WO 0204402	A	17-01-2002	AU 7102701 A	21-01-2002
			CA 2415468 A1	10-01-2003
			EP 1302458 A1	16-04-2003
			WO 0204402 A1	17-01-2002
<hr/>				